

A Counterion/Ligand-Tuned Chemo- and Enantioselective Copper-Catalyzed Intermolecular Radical 1,2-Carboamination of Alkenes

Xian-Yan Cheng,[§] Yu-Feng Zhang,^{*,§} Jia-Huan Wang, Qiang-Shuai Gu, Zhong-Liang Li,^{*} and Xin-Yuan Liu^{*}Cite This: <https://doi.org/10.1021/jacs.2c08035>

Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: The copper-catalyzed enantioselective intermolecular radical 1,2-carboamination of alkenes with readily accessible alkyl halides is an appealing strategy for producing chiral amine scaffolds. The challenge arises from the easily occurring atom transfer radical addition between alkyl halides and alkenes and the issue of enantiocontrol. We herein describe a radical alkene 1,2-carboamination with sulfoximines in a highly chemo- and enantioselective manner. The key to the success of this process is the conceptual design of a counterion/highly sterically demanded ligand coeffect to promote the ligand exchange of copper(I) with sulfoximines and forge chiral C–N bonds between alkyl radicals and the chiral copper(II) complex. The reaction covers alkenes bearing distinct electronic properties, such as aryl-, heteroaryl-, carbonyl-, and aminocarbonyl-substituted ones, and various radical precursors, including alkyl chlorides, bromides, iodides, and the CF₃ source. Facile transformations deliver many chiral amine building blocks of interest in organic synthesis and related areas.




INTRODUCTION

Alkenes are ubiquitous feedstocks in all aspects of chemistry and are readily available starting materials for versatile transformations. The intermolecular enantioselective 1,2-difunctionalization of alkenes represents a powerful tool for upgrading the feedstocks into enantioenriched complex molecules since it allows simultaneous installation of two vicinal new bonds.¹ Given the significance of chiral amines in pharmaceuticals and organic synthesis,² the intermolecular enantioselective 1,2-carboamination of alkenes has attracted great interest (Scheme 1A). Particularly, the precious transition metal catalysis has been developed to achieve such transformations, where the reaction initiation and enantiocontrol are generally achieved via the alkene carbometallation, a strategy pioneered by Cramer, Rovis, and others.^{3,4} In comparison, the radical-mediated intermolecular alkene 1,2-carboamination represents an important complementary approach because radicals have a high propensity to unsaturated bonds,⁵ thus avoiding the utilization of precious transition metals for alkene activation.⁶ Moreover, the abundance and easy availability of radical precursors further enriched the product diversity. As such, great progress has been made in developing the racemic intermolecular radical 1,2-carboamination of alkenes by harnessing the advantages of radical reactions.^{6,7} Among them, three pathways account for the C–N bond formation after the addition of carbon-centered radical to alkenes: (i) the direct radical–radical cross-coupling

or radical addition to unsaturated bonds;^{7a–c} (ii) the radical–cation crossover followed by nucleophilic addition;^{7d–i} and (iii) the 3d transition metal-mediated bond formation process (Scheme 1B).^{7j–n} Notably, the lack of polar functional groups nearby the alkyl radical center renders the enantiocontrol over the radical extremely difficult through the first two pathways.^{8,9} Although the last pathway provides a ready mechanism for enantiocontrol as pioneered by Fu and Peters,^{10,11} it has been seldom used in the intermolecular enantioselective alkene carboamination.¹² Only until very recently, Bao and others have realized a unique class of asymmetric alkene 1,2-carboazidation with TMSN₃, where an aryl substituent on the alkenes is necessary to deliver high enantioselectivity.¹³ Notably, the early-stage conversion of the chiral alkylazides to protected amines is generally required in a multiple-step transformation of the obtained chiral azide compounds.¹³ Therefore, the development of an intermolecular enantioselective 1,2-carboamination of diverse alkenes bearing distinct electronic properties with readily accessible radical precursors

Received: July 29, 2022

Scheme 1. Enantioselective Intermolecular Radical 1,2-Carboamination of Alkenes



and new amine nucleophiles that tolerate multiple-step synthetic operations is highly desirable.


We have been investigating copper-catalyzed asymmetric radical reactions¹⁴ and recently reported an enantioselective C(sp³)–N cross-coupling of alkyl halides with sulfoximines through the interaction of alkyl radicals with L*Cu^{II}Nu.¹⁵ Given the ready availability of alkyl halides, we wondered whether a three-component enantioselective alkene 1,2-carboamination can be achieved by merging our enantioselective C–N formation with the radical alkene addition process using sulfoximines as the amine sources.¹⁶ However, the major challenge would be the inherently facile atom transfer radical addition (ATRA)¹⁷ between alkyl halides and alkenes, which is one of the most prevailing strategies for alkene difunctionalization (Kharasch addition, the formation of ATRA product IV in Scheme 1C).¹⁸ Our initial attempts with styrene, sulfoximine **2a**, and CCl₄ showed that CuI and the chiral *N,N,P*-ligand **L1**^{15,19} could only afford the ATRA product (Scheme S1 in the Supporting Information). A further control experiment revealed that the ATRA product was also generated in a similar yield in the absence of sulfoximine **2a** (Scheme S1 in the Supporting Information). These results demonstrated that the ligand exchange with sulfoximine is so slow that the desired reaction pathway cannot compete with the ATRA process (Scheme 1C, pathway a versus c). To this end, we disclose a copper-catalyzed intermolecular radical 1,2-carboamination of alkenes with alkyl halides and sulfoximines with high chemo- and enantioselectivity. The key to the success of this process is the conceptual design of a combined counterion/chiral ligand effect to promote the desired process and suppress the easily occurring ATRA reaction: the counterion plays a pivotal role in accelerating the ligand exchange of sulfoximines with the copper catalyst; the highly sterically demanded chiral *N,N,P*-

ligand forges the desired chiral C–N bond instead of the C–X bond in the last step (Scheme 1C, pathway a versus b). The reaction exhibits a remarkably broad substrate scope, covering aryl-, heteroaryl-, carbonyl-, and aminocarbonyl-substituted alkenes bearing distinct electronic properties as well as diverse radical precursors, such as alkyl chlorides, bromides, iodides, and the CF₃ source (Scheme 1C). Owing to the broad scope of alkenes and radical precursors, the straightforward transformation enables it to be a powerful tool in generating diverse chiral amine building blocks.

RESULTS AND DISCUSSION

Reaction Development. With the preliminary reaction outcome in hand (Scheme S1 and Table 1, entries 1 and 2),

Table 1. Investigation of Counterions and Ligands for Model Reaction^a



L1, Ar = 3,5-(3,5-Me₂C₆H₃)₂C₆H₃ **L5**, Ar = 3,5-(3,5-Me₂C₆H₃)₂C₆H₃ X-ray structure of **4**

L2, Ar = Ph


L3, Ar = 3,5-Ph₂C₆H₃

L4, Ar = 3,5-(3,5-^tBu₂C₆H₃)₂C₆H₃

entry	[Cu]	L	additive	y. of 4 (%)	y. of 4' (%)	ee of 4 (%)
1 ^b	CuI	L1		trace	88	
2 ^c	CuI	L1		0	88	
3	CuI	L1	AcOH	45	10	96
4	CuI	L1	PivOH	89	3	97
5	CuI	L1	BzOH	45	15	95
6	CuI	L1	TcH	40	10	96
7	CuOAc	L1		43	0	98
8	CuTc	L1		90	0	96
9 ^d	CuTc	L1		85	0	96
10	CuTc	L2		2	24	59
11	CuTc	L3		10	35	82
12	CuTc	L4		11	0	96
13	CuTc	L5		94	0	–95

^aReaction conditions: **1a** (1.2 equiv), **2a** (0.05 mmol), **3a** (1.2 equiv), [Cu] (10 mol %), L (12 mol %), and Cs₂CO₃ (4.0 equiv) in Et₂O (0.5 mL) at room temperature for 5 days under argon; yields (y.) were based on ¹H NMR analysis of the crude product using 1,3,5-trimethoxybenzene as an internal standard; ee values were determined by HPLC analysis; all reactions were repeated twice. ^bThe ee values of **4'** were less than 5% during the whole optimization process. ^c**2a** was removed. ^dConducted at 40 °C for 1 day. PivOH, pivalic acid; Bz, benzoyl; and Tc, thiophene-2-carboxylate.

our endeavor toward the promotion of the desired pathway focused on accelerating the ligand exchange of sulfoximines with the copper catalyst. Inspired by the unique role of carboxylate anions in transition metal-catalyzed C–H activation via concerted metalation deprotonation (CMD),²⁰ we added carboxylic acid as an additive to assist the ligand exchange process. As such, several carboxylic acids were

Table 2. Scope of Sulfoximines and Radical Precursors^a

^aReaction conditions: **1** (1.2 equiv), **2** (0.20 mmol), **3** (alkyl chloride, 1.2 equiv), CuTc (10 mol %), **L1** (12 mol %), and Cs₂CO₃ (4.0 equiv) in Et₂O (2.0 mL) at room temperature for 5 days under argon; yields were isolated ones; ee values were determined by HPLC analysis. ^bCu(HFac)₂ was used. ^cAlkyl iodide was used. ^d1-Trifluoromethyl-1,2-benziodoxol-3(1H)-one (Togni reagent) was used. ^eHFac, hexafluoroacetylacetate.


investigated in the reaction of styrene **1a**, sulfoximine **2a**, and CCl₄ **3a**. We are pleased to find a dramatic suppression of the ATRA product **4'** and the formation of the desired carboamination product **4** (Table 1, entries 3–6). The pivalic acid, performing an essential role in the CMD process,^{20d} gave the best result with the formation of a trace amount of **4'** (Table 1, entry 4). We theorized that the low chemoselectivity of **4** with other carboxylic acids was possibly due to the equilibrium concentrations of copper iodide and carboxylate, which led to different reaction outcomes (Table 1, entries 3–6). Thus, we are encouraged to use pure copper(I) carboxylate as the catalyst to suppress the ATRA process. Gratifyingly, employing the commercially available CuOAc or CuTc salt significantly improved the chemoselectivity and the latter gave an excellent yield of **4** (Table 1, entries 7 and 8). Enhancing the reaction temperature to 40 °C could greatly increase the reaction rate to deliver **4** in a high yield without affecting the ee after 1 day (Table 1, entry 9). Further ligand investigation showed the steric effect on the *P*-aryl ring was also crucial for the chemoselectivity: while ligands with the less sterically hindered substituents (Dixon's ligand¹⁹ **L2** and **L3**) afforded low chemoselectivity and efficiency, the one with a bulkier substituent (**L4**) delivered **4** in only 11% yield (Table 1, entries 10–12). We reasoned that the interaction of the prochiral alkyl radical intermediate **I** with the Cu^{II} complex **II** to forge the C–N and C–X bond was greatly affected by the ligand architecture (Scheme 1C, pathway a versus b). The

small sterically hindered ligand was unfavorable for the C–N formation but preferred the C–X formation in the last step. The large sterically hindered ligand would prevent the interaction of intermediates **I** with **II** and be deleterious to the whole transformation. Considering the operational simplicity, we finally identified the optimal conditions as follows: **1a** (1.2 equiv), **2a** (1.0 equiv), **3a** (1.2 equiv), CuTc (10 mol %), **L1** (12 mol %), and Cs₂CO₃ (4.0 equiv) in Et₂O at room temperature, affording **4** in 90% NMR yield with 96% ee (Table 1, entry 8). The absolute configuration of **4** was determined to be *S* by X-ray analysis (Table 1 and Figure S1 in the Supporting Information) and those of other products were assigned by analogy. Replacing **L1** with its pseudoeantiomer **L5** afforded the enantiomer of **4** in 94% yield with 95% ee (Table 1, entry 13).

Substrate Scope. With the established optimal conditions in hand, we first evaluated the scope of sulfoximines (Table 2). The result showed that the reaction efficiency was influenced by the electronic effect on the aryl ring of sulfoximines: while the electron-donating or neutral substituents gave the desired carboamination products (**4**–**7**) with satisfactory yields and ee, the electron-withdrawing substituent was detrimental to the reaction efficiency (**8**). We next investigated the scope of radical precursors. Alkyl chlorides, such as trichlorotrifluoroethane and ethyl trichloroacetate, reacted well to result in **9**–**11** with excellent ee. A range of alkyl bromides that is α to ester or amide functional groups was amenable to the reactions, providing **12**–**16** with good ee. Furthermore, the fluoroalkyl iodide also proved to be suitable for the reaction to afford **17** in 91% yield with 87% ee. Most significantly, the Togni reagent reacted smoothly to furnish the pharmaceutically relevant β -trifluoromethylated amine **18** in 94% yield with 97% ee. However, the reaction could not be initiated with primary alkyl halides, such as ethyl chloroacetate and ethyl bromoacetate.

We then investigated the scope of alkenes for the asymmetric 1,2-carboamination. As shown in Table 3, the styrene-type alkenes possessing diverse electron-donating and -withdrawing substituents at different (*ortho*, *meta*, or *para*) positions of the phenyl rings reacted smoothly to afford **19**–**35** in 70–98% yields with up to 99% ee. Importantly, a gamut of reactive functionalities, such as acetyl (**22**), thioether (**23**), halogen (**26**–**28**), nitrile (**30**), formyl (**31** and **32**), ketone (**33**), ester (**34**), and even nitro (**35**) groups, were well tolerated under the standard conditions. In addition, the phenyl ring having mono- and disubstituted conjugated groups (phenyl, terminal, and internal alkene, alkyne) was compatible with the reaction conditions to deliver **36**–**40** in moderate to good yields with high enantioselectivities. Alkenes bearing polycyclic aryl rings were also suitable for the reactions, giving rise to **41**–**47** in good yields with 87 to >99% ee. More importantly, alkenes containing many medically relevant heterocycles, such as pyridine, quinoline, thiophene, and benzo[*b*]thiophene, were viable substrates to furnish the desired products **48**–**51** with up to 99% ee.

Enantioenriched α -amino carbonyl compounds are a class of privileged motifs in many pharmaceutically active agents and valuable intermediates in organic synthesis.²¹ To further strengthen the utility of the strategy, we next switched our attention to the scope of other carbonyl-substituted alkenes, which have not been utilized in the reported enantioselective radical alkene difunctionalization.^{12b,13} We found that an array of carbonyl-substituted alkenes was suitable for the reaction to

Table 3. Scope of Styrene-Type Alkenes^a


^aReaction conditions: **1** (1.2 equiv), **2a** (0.20 mmol), **3a** (1.2 equiv), CuTc (10 mol %), L1 (12 mol %), and Cs₂CO₃ (4.0 equiv) in Et₂O (2.0 mL) at room temperature for 5 days. ^b4-Cyclohexyl-substituted diaryl sulfoximine (**2b**) was used; yields were isolated ones; ee values were determined by HPLC analysis.

provide enantioenriched α -amino ketones **52–56** in moderate to good yields with 81–96% ee under slightly modified conditions (Table 4). Both aryl and alkyl ketones were compatible with the reaction conditions. Notably, the nitrile-substituted alkene was also a viable substrate, resulting in the chiral α -amino nitrile **57**, albeit with a moderate ee. More significantly, a panel of amide-substituted alkenes was well accommodated, leading to enantioenriched α -amino amides **58–61** with 81–92% ee. As for the carbonyl-substituted alkenes, the high reaction efficiency and enantioselectivity might be attributed to the coordination of the copper catalyst with the substrates.²² Collectively, this strategy not only achieved the enantioselective radical 1,2-difunctionalization of carbonyl-substituted alkenes for the first time but also provided diverse enantioenriched γ -functionalized α -amino carbonyl compounds.

Synthetic Utility. To show the preparative utility of this method, a gram-scale reaction was first carried out to furnish the desired product **20** without a decrease in efficiency (Scheme 2A). The most important application is that it provides a versatile platform for expedient synthesis of diverse chiral amine building blocks. To showcase this potential, the enantioenriched sulfoximine **18** was successfully converted to the chiral β -trifluoromethylated amine derivative **62** using a sequential Na/naphthalene reduction/acidic hydrolysis/protection protocol (Scheme 2B). The facile functionality transformation/reduction of trichloromethylated product **20**

afforded an γ -amino aldehyde **63** and further delivered **64** via a Wittig reaction, which was converted to an enantioenriched δ -alkenyl amine building block **65** (Scheme 2C). Moreover, the dichloromethylated product **11** also underwent a reduction process to furnish a γ -amino acid derivative **66**. The direct deprotection or reduction/deprotection of **66** resulted in the enantioenriched γ -lactam (**67**) and 1,4-amino alcohol (**68**) building blocks. Notably, no apparent loss of enantiopurity was observed during all these transformations. Notably, the sulfoximine moieties are proved to be stable protecting amine functionalities and could tolerate multistep synthesis, as demonstrated in the synthesis of **65–68**.

Mechanistic Consideration. To gain insights into the reaction mechanism, several control experiments were conducted. A radical trap experiment with butylhydroxytoluene (BHT) showed that the reaction was completely inhibited and the BHT-trapped product **69** was detected (Scheme 3A). The radical clock reaction with **70** bearing an α cyclopropyl substituent delivered the radical addition/ring-opening/C–N formation product **71** in 12% yield under the standard reaction conditions. The combined experiments indicated the involvement of a radical process (Scheme 3B). Notably, no coupling product between **4'** and **2a** was formed when the ATRA side product **4'** was exposed to the reaction of **1b** and only **19** was afforded. In comparison, both **4** and **19** were obtained when the corresponding alkenes **1a** and **1b** were simultaneously exposed to the standard reaction conditions

Table 4. Scope of Electron-Deficient Alkenes^a

^aReaction conditions: **1** (2.0 equiv), **2** (0.20 mmol), **3** (alkyl bromides, 2.0 equiv), Cu(HFac)₂ (10 mol %), L1 (12 mol %), and Cs₂CO₃ (4.0 equiv) in Et₂O (2 mL) at 0 °C for 5 days. ^b**1** (1.5 equiv), **2** (0.20 mmol), **3** (alkyl bromides, 1.5 equiv) were used. ^cL4 (12 mol %) were used. ^dConducted at room temperature; yields were isolated ones; ee values were determined by HPLC analysis.

(Schemes 3C and S2 in the Supporting Information). Meanwhile, the direct coupling of **4'** and **2a** did not occur under the standard conditions as well (Scheme S2 in the Supporting Information). Collectively, these experiments together with the formation of the carboamination product **18** excluded the possibility of a stepwise atom transfer radical addition/cross-coupling process (Scheme 3C and Table 2).

Based on the above experiments and literature reports,^{14,15} we proposed a possible mechanism as depicted in Scheme 3D. First, the copper(I) salt reacted with ligand L and base to generate a complex A. Without the facilitation of the counterion, complex A would directly react with the radical precursors **3** to generate the ATRA side product IV via the

interaction of I and III (the right-hand catalytic cycle). The addition of a carboxylate anion would switch the reaction pathway through the formation of a copper(I) carboxylate B, which underwent a fast ligand exchange with sulfoximines **2** to afford a copper(I)-sulfoximinato complex C (the left-hand catalytic cycle) via a CMD-type process.²⁰ This intermediate then reduced the radical precursors **3**, leading to the copper(II)-sulfoximinato complex II and an R¹ radical. The R¹ radical underwent a facile addition to alkenes **1** and provided the prochiral alkyl radical I. Next, radical I interacted with complex II to deliver the desired carboamination products **4-61** and regenerated the copper(I) species A in the presence of the highly sterically demanded ligand L1. As mentioned in the reaction development section, both the less and more sterically hindered ligands are harmful to the desired pathway by either forming the side product IV (pathway b in Scheme 1C) or retarding the interaction of I and II.

CONCLUSIONS


In summary, we have developed a copper-catalyzed intermolecular enantioselective radical 1,2-carboamination of alkenes from readily available alkyl halides and sulfoximines by successfully merging the radical addition to alkenes and enantioselective radical C–N formation. The employment of a carboxylate counterion and the highly sterically demanded *N,N,P*-ligand is vital for the high chemo- and enantioselectivity. More significantly, the alkene scope is very broad, covering not only aryl and heteroaryl alkenes but also carbonyl-substituted alkenes that have not been disclosed in reported asymmetric radical alkene 1,2-difunctionalization. A wide array of alkyl halide radical precursors is easily accommodated into alkenes under ambient conditions. This approach provides a versatile platform for the synthesis of chiral amines as well as other synthetically valuable chiral building blocks.

ASSOCIATED CONTENT


Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.2c08035>.

Scheme 2. Synthetic Utility



Scheme 3. Mechanistic Investigation



Experimental procedures, characterization of compounds, initial test on copper-catalyzed enantioselective 1,2-carboamination, control experiments to exclude the stepwise ATRA/cross-coupling process, and crystallographic data of **4** ([PDF](#))

Accession Codes

CCDC 2192979 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Authors

Yu-Feng Zhang – Shenzhen Grubbs Institute and Department of Chemistry, Guangdong Provincial Key Laboratory of Catalysis, Southern University of Science and Technology, Shenzhen 518055, China; Email: zhangyf3@sustech.edu.cn

Zhong-Liang Li – Academy for Advanced Interdisciplinary Studies and Department of Chemistry, Southern University of Science and Technology, Shenzhen 518055, China;
Email: lizl@sustech.edu.cn

Xin-Yuan Liu — Shenzhen Grubbs Institute and Department of Chemistry, Guangdong Provincial Key Laboratory of Catalysis, Southern University of Science and Technology, Shenzhen 518055, China; orcid.org/0000-0002-6978-6465; Email: liuxy3@sustech.edu.cn


Authors

Xian-Yan Cheng – *Shenzhen Grubbs Institute and
Department of Chemistry, Guangdong Provincial Key*

Laboratory of Catalysis, Southern University of Science and
Technology, Shenzhen 518055, China

Jia-Huan Wang – Shenzhen Grubbs Institute and Department of Chemistry, Guangdong Provincial Key Laboratory of Catalysis, Southern University of Science and Technology, Shenzhen 518055, China

Qiang-Shuai Gu – Academy for Advanced Interdisciplinary Studies and Department of Chemistry, Southern University of Science and Technology, Shenzhen 518055, China;

 orcid.org/0000-0002-3840-425X

Complete contact information is available at:
<https://pubs.acs.org/10.1021/jacs.2c08035>

Author Contributions

[§]X.-Y.C. and Y.-F.Z. contributed equally.

Author Contributions

The manuscript was written through the contributions of all authors.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial support from the National Key R&D Program of China (Nos. 2021YFF0701604 and 2021YFF0701704), National Natural Science Foundation of China (Nos. 22025103 and 21831002), Guangdong Innovative Program (No. 2019BT02Y335), Guangdong Provincial Key Laboratory of Catalysis (No. 2020B121201002), Guangdong Basic and Applied Basic Research Foundation (No. 2021A1515110093), and Shenzhen Special Funds (No. JCYJ20200109141001789) is acknowledged. The authors

acknowledge the assistance of SUSTech Core Research Facilities. This study is dedicated to Professor Pei-Qiang Huang on the occasion of his 60th birthday.

REFERENCES

- (1) For selected reviews on intermolecular alkene 1,2-difunctionalization, see: (a) McDonald, R. I.; Liu, G.; Stahl, S. S. Palladium(II)-Catalyzed Alkene Functionalization via Nucleopalladation: Stereochemical Pathways and Enantioselective Catalytic Applications. *Chem. Rev.* **2011**, *111*, 2981–3019. (b) Shimizu, Y.; Kanai, M. Recent Progress in Copper-Catalyzed Difunctionalization of Unactivated Carbon–Carbon Multiple Bonds. *Tetrahedron Lett.* **2014**, *55*, 3727–3737. (c) Lee, J. H.; Choi, S.; Hong, K. B. Alkene Difunctionalization Using Hypervalent Iodine Reagents: Progress and Developments in the Past Ten Years. *Molecules* **2019**, *24*, 2634. (d) Derosa, J.; Apolinar, O.; Kang, T.; Tran, V. T.; Engle, K. M. Recent Developments in Nickel-Catalyzed Intermolecular Dicarbofunctionalization of Alkenes. *Chem. Sci.* **2020**, *11*, 4287–4296. (e) Bhunia, A.; Studer, A. Recent Advances in Radical Chemistry Proceeding through Pro-aromatic Radicals. *Chem* **2021**, *7*, 2060–2100.
- (2) (a) Nugent, T. C. *Chiral Amine Synthesis: Methods, Developments and Applications*; Wiley: Weinheim, 2010; pp 1–506. (b) Ricci, A.; Bernardi, L. *Methodologies in Amine Synthesis: Challenges and Applications*; Wiley: Weinheim, 2021; pp 1–460.
- (3) For leading examples, see: (a) Hyster, T. K.; Knörr, L.; Ward, T. R.; Rovis, T. Biotinylated Rh(III) Complexes in Engineered Streptavidin for Accelerated Asymmetric C–H Activation. *Science* **2012**, *338*, 500–503. (b) Ye, B.; Cramer, N. Chiral Cyclopentadienyl Ligands as Stereocontrolling Element in Asymmetric C–H Functionalization. *Science* **2012**, *338*, 504–506. For other selected examples, see: (c) Chen, S.-S.; Wu, M.-S.; Han, Z.-Y. Palladium-Catalyzed Cascade sp^2 C–H Functionalization/Intramolecular Asymmetric Allylation: From Aryl Ureas and 1,3-Dienes to Chiral Indolines. *Angew. Chem., Int. Ed.* **2017**, *56*, 6641–6645. (d) Teng, H.-L.; Luo, Y.; Nishiura, M.; Hou, Z. Diastereodivergent Asymmetric Carboamination/Annulation of Cyclopropenes with Aminoalkenes by Chiral Lanthanum Catalysts. *J. Am. Chem. Soc.* **2017**, *139*, 16506–16509. (e) Tao, M.; Tu, Y.; Liu, Y.; Wu, H.; Liu, L.; Zhang, J. Pd/Xiang-Phos-Catalyzed Enantioselective Intermolecular Carboheterofunctionalization under Mild Conditions. *Chem. Sci.* **2020**, *11*, 6283–6288. (f) Cui, W.-J.; Wu, Z.-J.; Gu, Q.; You, S.-L. Divergent Synthesis of Tunable Cyclopentadienyl Ligands and Their Application in Rh-Catalyzed Enantioselective Synthesis of Isoindolinone. *J. Am. Chem. Soc.* **2020**, *142*, 7379–7385. (g) Mi, R.; Zhang, X.; Wang, J.; Chen, H.; Lan, Y.; Wang, F.; Li, X. Rhodium-Catalyzed Regio-, Diastereo-, and Enantioselective Three-Component Carboamination of Dienes via C–H Activation. *ACS Catal.* **2021**, *11*, 6692–6697. (h) Wu, L.; Xu, H.; Gao, H.; Li, L.; Chen, W.; Zhou, Z.; Yi, W. Chiral Allylic Amine Synthesis Enabled by the Enantioselective $Cp^*Rh(III)$ -Catalyzed Carboaminations of 1,3-Dienes. *ACS Catal.* **2021**, *11*, 2279–2287. For one example using chiral cobalt catalysis, see: (i) Ozols, K.; Jang, Y.-S.; Cramer, N. Chiral Cyclopentadienyl Cobalt(III) Complexes Enable Highly Enantioselective 3d-Metal-Catalyzed C–H Functionalizations. *J. Am. Chem. Soc.* **2019**, *141*, 5675–5680.
- (4) For selected reviews on asymmetric intramolecular 1,2-carboamination, see: (a) Schultz, D. M.; Wolfe, J. P. Recent Developments in Palladium-Catalyzed Alkene Aminoarylation Reactions for the Synthesis of Nitrogen Heterocycles. *Synthesis* **2012**, *44*, 351–361. (b) White, D. R.; Bornowski, E. C.; Wolfe, J. P. Pd-Catalyzed C–C, C–N, and C–O Bond-Forming Difunctionalization Reactions of Alkenes Bearing Tethered Aryl/Alkenyl Triflates. *Isr. J. Chem.* **2020**, *60*, 259–267. For selected examples, see: (c) Mai, D. N.; Wolfe, J. P. Asymmetric Palladium-Catalyzed Carboamination Reactions for the Synthesis of Enantiomerically Enriched 2-(Arylmethyl)- and 2-(Alkenylmethyl)pyrrolidines. *J. Am. Chem. Soc.* **2010**, *132*, 12157–12159. (d) Hopkins, B. A.; Wolfe, J. P. Synthesis of Enantiomerically Enriched Imidazolidin-2-Ones through Asymmetric Palladium-Catalyzed Alkene Carboamination Reactions. *Angew. Chem., Int. Ed.* **2012**, *51*, 9886–9890. (e) White, D. R.; Hutt, J. T.; Wolfe, J. P. Asymmetric Pd-Catalyzed Alkene Carboamination Reactions for the Synthesis of 2-Aminoindane Derivatives. *J. Am. Chem. Soc.* **2015**, *137*, 11246–11249.
- (5) Giese, B. Formation of CC Bonds by Addition of Free Radicals to Alkenes. *Angew. Chem., Int. Ed. Engl.* **1983**, *22*, 753–764.
- (6) For selected reviews on intermolecular radical 1,2-carboamination of alkenes, see: (a) Jiang, H.; Studer, A. Intermolecular Radical Carboamination of Alkenes. *Chem. Soc. Rev.* **2020**, *49*, 1790–1811. (b) Koike, T.; Akita, M. New Horizons of Photocatalytic Fluoromethylative Difunctionalization of Alkenes. *Chem* **2018**, *4*, 409–437. (c) Chen, X.; Xiao, F.; He, W.-M. Recent Developments in the Difunctionalization of Alkenes with C–N Bond Formation. *Org. Chem. Front.* **2021**, *8*, 5206–5228. (d) Wu, Z.; Hu, M.; Li, J.; Wu, W.; Jiang, H. Recent Advances in Aminative Difunctionalization of Alkenes. *Org. Biomol. Chem.* **2021**, *19*, 3036–3054.
- (7) For selected examples of intermolecular radical 1,2-carboamination of alkenes: (a) Patra, T.; Bellotti, P.; Strieth-Kalthoff, F.; Glorius, F. Photosensitized Intermolecular Carboamination of Alkenes through the Persistent Radical Effect. *Angew. Chem., Int. Ed.* **2020**, *59*, 3172–3177. (b) Kindt, S.; Wicht, K.; Heinrich, M. R. Base-Induced Radical Carboamination of Nonactivated Alkenes with Aryldiazonium Salts. *Org. Lett.* **2015**, *17*, 6122–6125. (c) Blank, O.; Heinrich, M. R. Carbodiazenylation of Olefins by Radical Iodine Transfer and Addition to Arenediazonium Salts. *Eur. J. Org. Chem.* **2006**, *2006*, 4331–4334. (d) Fumagalli, G.; Boyd, S.; Greaney, M. F. Oxyarylation and Aminoarylation of Styrenes Using Photoredox Catalysis. *Org. Lett.* **2013**, *15*, 4398–4401. (e) Yasu, Y.; Koike, T.; Akita, M. Intermolecular Aminotrifluoromethylation of Alkenes by Visible-Light-Driven Photoredox Catalysis. *Org. Lett.* **2013**, *15*, 2136–2139. (f) Prasad Hari, D.; Hering, T.; König, B. The Photoredox-Catalyzed Meerwein Addition Reaction: Intermolecular Amino-Arylation of Alkenes. *Angew. Chem., Int. Ed.* **2014**, *53*, 725–728. (g) Liu, Y.-Y.; Yang, X.-H.; Song, R.-J.; Luo, S.; Li, J.-H. Oxidative 1,2-Carboamination of Alkenes with Alkyl Nitriles and Amines toward γ -Amino Alkyl Nitriles. *Nat. Commun.* **2017**, *8*, 14720. (h) Qian, B.; Chen, S.; Wang, T.; Zhang, X.; Bao, H. Iron-Catalyzed Carboamination of Olefins: Synthesis of Amines and Disubstituted β -Amino Acids. *J. Am. Chem. Soc.* **2017**, *139*, 13076–13082. (i) Gockel, S. N.; Buchanan, T. L.; Hull, K. L. Cu-Catalyzed Three-Component Carboamination of Alkenes. *J. Am. Chem. Soc.* **2018**, *140*, 58–61. (j) Xiong, Y.; Ma, X.; Zhang, G. Copper-Catalyzed Intermolecular Carboamination of Alkenes Induced by Visible Light. *Org. Lett.* **2019**, *21*, 1699–1703. (k) Gockel, S. N.; Lee, S.; Gay, B. L.; Hull, K. L. Oxidative Three-Component Carboamination of Vinylarenes with Alkylboronic Acids. *ACS Catal.* **2021**, *11*, 5166–5171. (l) Lu, D.; Li, Y.; Wang, P.; Wang, Z.; Yang, D.; Gong, Y. Cu-Catalyzed $C(sp^3)$ –N Coupling and Alkene Carboamination Enabled by Ligand-Promoted Selective Hydrazine Transfer to Alkyl Radicals. *ACS Catal.* **2022**, *12*, 3269–3278. (m) Ma, D.; Kong, D.; Wu, P.; Tu, Y.; Shi, P.; Wang, X.; Bolm, C. Introduction of Lipophilic Side Chains to NH-Sulfoximines by Palladium Catalysis Under Blue Light Irradiation. *Org. Lett.* **2022**, *24*, 2238–2241. (n) Kennedy-Ellis, J. J.; Boldt, E. D.; Chemler, S. R. Synthesis of Benzylureas and Related Amine Derivatives via Copper-Catalyzed Three-Component Carboamination of Styrenes. *Org. Lett.* **2020**, *22*, 8365–8369.
- (8) The asymmetric radical reactions of direct coupling or addition strategy necessitates a polar functional group nearby the radical center, for representative reviews or examples, see: (a) Sibi, M. P.; Manyem, S.; Zimmerman, J. Enantioselective Radical Processes. *Chem. Rev.* **2003**, *103*, 3263–3296. (b) Nicewicz, D. A.; MacMillan, D. W. C. Merging Photoredox Catalysis with Organocatalysis: The Direct Asymmetric Alkylation of Aldehydes. *Science* **2008**, *322*, 77–80. (c) Du, J.; Skubi, K. L.; Schultz, D. M.; Yoon, T. P. A Dual-Catalysis Approach to Enantioselective [2 + 2] Photocycloadditions Using Visible Light. *Science* **2014**, *344*, 392–396. (d) Hashimoto, T.; Kawamata, Y.; Maruoka, K. An Organic Thiyl Radical Catalyst for Enantioselective Cyclization. *Nat. Chem.* **2014**, *6*, 702–705.

(e) Brimioulle, R.; Lenhart, D.; Maturi, M. M.; Bach, T. Enantioselective Catalysis of Photochemical Reactions. *Angew. Chem., Int. Ed.* **2015**, *54*, 3872–3890. (f) Murphy, J. J.; Bastida, D.; Paria, S.; Fagnoni, M.; Melchiorre, P. Asymmetric Catalytic Formation of Quaternary Carbons by Iminium Ion Trapping of Radicals. *Nature* **2016**, *532*, 218–222. (g) Kern, N.; Plesniak, M. P.; McDouall, J. J. W.; Procter, D. J. Enantioselective Cyclizations and Cyclization Cascades of Samarium Ketyl Radicals. *Nat. Chem.* **2017**, *9*, 1198–1204. (h) Proctor, R. S. J.; Davis, H. J.; Phipps, R. J. Catalytic Enantioselective Minisci-Type Addition to Heteroarenes. *Science* **2018**, *360*, 419–422. (i) Biegasiewicz, K. F.; Cooper, S. J.; Gao, X.; Oblinsky, D. G.; Kim, J. H.; Garfinkle, S. E.; Joyce, L. A.; Sandoval, B. A.; Scholes, G. D.; Hyster, T. K. Photoexcitation of Flavoenzymes Enables a Stereoselective Radical Cyclization. *Science* **2019**, *364*, 1166–1169.

(9) For one example of the efficient enantiocontrol over purely carbocation intermediate without an adjacent polar functional group, see: Wendlandt, A. E.; Vangal, P.; Jacobsen, E. N. Quaternary Stereocentres via an Enantioconvergent Catalytic S_N1 Reaction. *Nature* **2018**, *556*, 447–451.

(10) (a) Kainz, Q. M.; Matier, C. D.; Bartoszewicz, A.; Zultanski, S. L.; Peters, J. C.; Fu, G. C. Asymmetric Copper-Catalyzed C–N Cross-Couplings Induced by Visible Light. *Science* **2016**, *351*, 681–684. (b) Chen, C.; Peters, J. C.; Fu, G. C. Photoinduced Copper-Catalyzed Asymmetric Amidation via Ligand Cooperativity. *Nature* **2021**, *596*, 250–256. (c) Cho, H.; Suematsu, H.; Oyala, P. H.; Peters, J. C.; Fu, G. C. Photoinduced, Copper-Catalyzed Enantioconvergent Alkylations of Anilines by Racemic Tertiary Electrophiles: Synthesis and Mechanism. *J. Am. Chem. Soc.* **2022**, *144*, 4550–4558.

(11) (a) Wang, F.; Chen, P.; Liu, G. Copper-Catalyzed Radical Relay for Asymmetric Radical Transformations. *Acc. Chem. Res.* **2018**, *51*, 2036–2046. (b) Wang, Y.; Wen, X.; Cui, X.; Wojtas, L.; Zhang, X. P. Asymmetric Radical Cyclopropanation of Alkenes with In Situ-Generated Donor-Substituted Diazo Reagents via Co(II)-Based Metalloradical Catalysis. *J. Am. Chem. Soc.* **2017**, *139*, 1049–1052. (c) Yang, Y.; Cho, I.; Qi, X.; Liu, P.; Arnold, F. H. An Enzymatic Platform for the Asymmetric Amination of Primary, Secondary and Tertiary C(sp³)–H Bonds. *Nat. Chem.* **2019**, *11*, 987–993. (d) Nakafuku, K. M.; Zhang, Z.; Wappes, E. A.; Stateman, L. M.; Chen, A. D.; Nagib, D. A. Enantioselective Radical C–H Amination for the Synthesis of β -Amino Alcohols. *Nat. Chem.* **2020**, *12*, 697–704. (e) Zhou, Q.; Chin, M.; Fu, Y.; Liu, P.; Yang, Y. Stereodivergent Atom-Transfer Radical Cyclization by Engineered Cytochromes P450. *Science* **2021**, *374*, 1612–1616. (f) Huo, H.; Shen, X.; Wang, C.; Zhang, L.; Röse, P.; Chen, L.-A.; Harms, K.; Marsch, M.; Hilt, G.; Meggers, E. Asymmetric Photoredox Transition-Metal Catalysis Activated by Visible Light. *Nature* **2014**, *515*, 100–103. (g) Lang, K.; Hu, Y.; Lee, W.-C. C.; Zhang, X. P. Combined Radical and Ionic Approach for the Enantioselective Synthesis of β -Functionalized Amines from Alcohols. *Nat. Synth.* **2022**, *1*, 548–557. (h) Lang, K.; Li, C.; Kim, I.; Zhang, X. P. Enantioconvergent Amination of Racemic Tertiary C–H Bonds. *J. Am. Chem. Soc.* **2020**, *142*, 20902–20911. (i) Jin, L. M.; Xu, P.; Xie, J.; Zhang, X. P. Enantioselective Intermolecular Radical C–H Amination. *J. Am. Chem. Soc.* **2020**, *142*, 20828–20836. (j) Hu, Y.; Lang, K.; Li, C.; Gill, J. B.; Kim, I.; Lu, H.; Fields, K. B.; Marshall, M.; Cheng, Q.; Cui, X.; Wojtas, L.; Zhang, X. P. Enantioselective Radical Construction of 5-Membered Cyclic Sulfonamides by Metalloradical C–H Amination. *J. Am. Chem. Soc.* **2019**, *141*, 18160–18169. (k) Hu, Y.; Lang, K.; Tao, J.; Marshall, M. K.; Cheng, Q.; Cui, X.; Wojtas, L.; Zhang, X. P. Next-Generation D2-Symmetric Chiral Porphyrins for Cobalt(II)-Based Metalloradical Catalysis: Catalyst Engineering by Distal Bridging. *Angew. Chem., Int. Ed.* **2019**, *58*, 2670–2674.

(12) For selected reviews on the intramolecular version of asymmetric radical carboamination, see: (a) Chemler, S. R.; Karyakarte, S. D.; Khoder, Z. M. Stereoselective and Regioselective Synthesis of Heterocycles via Copper-Catalyzed Additions of Amine Derivatives and Alcohols to Alkenes. *J. Org. Chem.* **2017**, *82*, 11311–11325. (b) Li, Z.-L.; Fang, G.-C.; Gu, Q.-S.; Liu, X.-Y. Recent

Advances in Copper-Catalysed Radical-Involved Asymmetric 1,2-Difunctionalization of Alkenes. *Chem. Soc. Rev.* **2020**, *49*, 32–48. For selected examples on the intramolecular version, see: (c) Zeng, W.; Chemler, S. R. Copper(II)-Catalyzed Enantioselective Intramolecular Carboamination of Alkenes. *J. Am. Chem. Soc.* **2007**, *129*, 12948–12949. (d) Liwosz, T. W.; Chemler, S. R. Copper-Catalyzed Enantioselective Intramolecular Alkene Amination/Intermolecular Heck-Type Coupling Cascade. *J. Am. Chem. Soc.* **2012**, *134*, 2020–2023. (e) Lin, J.-S.; Dong, X.-Y.; Li, T.-T.; Jiang, N.-C.; Tan, B.; Liu, X.-Y. A Dual-Catalytic Strategy To Direct Asymmetric Radical Aminotrifluoromethylation of Alkenes. *J. Am. Chem. Soc.* **2016**, *138*, 9357–9360. (f) Bai, Z.; Zhang, H.; Wang, H.; Yu, H.; Chen, G.; He, G. Enantioselective Alkylamination of Unactivated Alkenes under Copper Catalysis. *J. Am. Chem. Soc.* **2021**, *143*, 1195–1202. (g) Riart-Ferrer, X.; Sang, P.; Tao, J.; Xu, H.; Jin, L.-M.; Lu, H.; Cui, X.; Wojtas, L.; Zhang, X. P. Metalloradical Activation of Carbonyl Azides for Enantioselective Radical Aziridination. *Chem* **2021**, *7*, 1120–1134. (h) Jiang, H.; Lang, K.; Lu, H.; Wojtas, L.; Zhang, X. P. Asymmetric Radical Bicyclization of Allyl Azidoformates via Cobalt(II)-Based Metalloradical Catalysis. *J. Am. Chem. Soc.* **2017**, *139*, 9164–9167. (i) Jin, L. M.; Xu, X.; Lu, H.; Cui, X.; Wojtas, L.; Zhang, X. P. Effective Synthesis of Chiral N-Fluoroaryl Aziridines through Enantioselective Aziridination of Alkenes with Fluoroaryl Azides. *Angew. Chem., Int. Ed.* **2013**, *52*, 5309–5313.

(13) (a) Ge, L.; Zhou, H.; Chiou, M.-F.; Jiang, H.; Jian, W.; Ye, C.; Li, X.; Zhu, X.; Xiong, H.; Li, Y.; Song, L.; Zhang, X.; Bao, H. Iron-Catalysed Asymmetric Carboazidation of Styrenes. *Nat. Catal.* **2021**, *4*, 28–35. (b) Liu, W.; Pu, M.; He, J.; Zhang, T.; Dong, S.; Liu, X.; Wu, Y.-D.; Feng, X. Iron-Catalyzed Enantioselective Radical Carboazidation and Diazidation of α,β -Unsaturated Carbonyl Compounds. *J. Am. Chem. Soc.* **2021**, *143*, 11856–11863. (c) Wu, L.; Zhang, Z.; Wu, D.; Wang, F.; Chen, P.; Lin, Z.; Liu, G. Anionic Bisoxazoline Ligands Enable Copper-Catalyzed Asymmetric Radical Azidation of Acrylamides. *Angew. Chem., Int. Ed.* **2021**, *60*, 6997–7001.

(14) For selected reviews on asymmetric radical reactions from our group, see: (a) Gu, Q.-S.; Li, Z.-L.; Liu, X.-Y. Copper(I)-Catalyzed Asymmetric Reactions Involving Radicals. *Acc. Chem. Res.* **2020**, *53*, 170–181. (b) Zhou, H.; Li, Z.-L.; Gu, Q.-S.; Liu, X.-Y. Ligand-Enabled Copper(I)-Catalyzed Asymmetric Radical C(sp³)–C Cross-Coupling Reactions. *ACS Catal.* **2021**, *11*, 7978–7986. (c) Dong, X.-Y.; Li, Z.-L.; Gu, Q.-S.; Liu, X.-Y. Ligand Development for Copper-Catalyzed Enantioconvergent Radical Cross-Coupling of Racemic Alkyl Halides. *J. Am. Chem. Soc.* **2022**, DOI: 10.1021/jacs.2c06718. For selected recent examples, see: (d) Dong, X.-Y.; Zhang, Y.-F.; Ma, C.-L.; Gu, Q.-S.; Wang, F.-L.; Li, Z.-L.; Jiang, S.-P.; Liu, X.-Y. A General Asymmetric Copper-Catalysed Sonogashira C(sp³)–C(sp) Coupling. *Nat. Chem.* **2019**, *11*, 1158–1166. (e) Jiang, S.-P.; Dong, X.-Y.; Gu, Q.-S.; Ye, L.; Li, Z.-L.; Liu, X.-Y. Copper-Catalyzed Enantioconvergent Radical Suzuki-Miyaura C(sp³)–C(sp²) Cross-Coupling. *J. Am. Chem. Soc.* **2020**, *142*, 19652–19659. (f) Su, X.-L.; Ye, L.; Chen, J.-J.; Liu, X.-D.; Jiang, S.-P.; Wang, F.-L.; Liu, L.; Yang, C.-J.; Chang, X.-Y.; Li, Z.-L.; Gu, Q.-S.; Liu, X.-Y. Copper-Catalyzed Enantioconvergent Cross-Coupling of Racemic Alkyl Bromides with Azole C(sp²)–H Bonds. *Angew. Chem., Int. Ed.* **2021**, *60*, 380–384. (g) Wang, P.-F.; Yu, J.; Guo, K.-X.; Jiang, S.-P.; Chen, J.-J.; Gu, Q.-S.; Liu, J.-R.; Hong, X.; Li, Z.-L.; Liu, X.-Y. Design of Hemilabile N,N,N-Ligands in Copper-Catalyzed Enantioconvergent Radical Cross-Coupling of Benzyl/Propargyl Halides with Alkenylboronate Esters. *J. Am. Chem. Soc.* **2022**, *144*, 6442–6452. (h) Wang, F.-L.; Yang, C.-J.; Liu, J.-R.; Yang, N.-Y.; Dong, X.-Y.; Jiang, R.-Q.; Chang, X.-Y.; Li, Z.-L.; Xu, G.-X.; Yuan, D.-L.; Zhang, Y.-S.; Gu, Q.-S.; Hong, X.; Liu, X.-Y. Mechanism-Based Ligand Design for Copper-Catalysed Enantioconvergent C(sp³)–C(sp) Cross-Coupling of Tertiary Electrophiles with Alkynes. *Nat. Chem.* **2022**, *14*, 949–957.

(15) Zhang, Y.-F.; Dong, X.-Y.; Cheng, J.-T.; Yang, N.-Y.; Wang, L.-L.; Wang, F.-L.; Luan, C.; Liu, J.; Li, Z.-L.; Gu, Q.-S.; Liu, X.-Y. Enantioconvergent Cu-Catalyzed Radical C–N Coupling of Racemic

Secondary Alkyl Halides to Access α -Chiral Primary Amines. *J. Am. Chem. Soc.* **2021**, *143*, 15413–15419.

(16) For one recent review, see: (a) Li, X.; Wang, C.; Jia, T. Recent Advances in *N*-Arylation of NH-Sulfoximines and Their Applications. *Chin. J. Org. Chem.* **2022**, *42*, 714–731. For selected examples, see: (b) Wang, L.; Priebbenow, D. L.; Zou, L.-H.; Bolm, C. The Copper-Catalyzed Oxidative *N*-Acylation of Sulfoximines. *Adv. Synth. Catal.* **2013**, *355*, 1490–1494. (c) Cheng, Y.; Dong, W.; Wang, L.; Parthasarathy, K.; Bolm, C. Iron-Catalyzed Hetero-Cross-Dehydrogenative Coupling Reactions of Sulfoximines with Diarylmethanes: A New Route to *N*-Alkylated Sulfoximines. *Org. Lett.* **2014**, *16*, 2000–2002. (d) Teng, F.; Sun, S.; Jiang, Y.; Yu, J.-T.; Cheng, J. Copper-Catalyzed Oxidative C(sp³)-H/N-H Coupling of Sulfoximines and Amides with Simple Alkanes via a Radical Process. *Chem. Commun.* **2015**, *51*, 5902–5905.

(17) Kharasch, M. S.; Jensen, E. V.; Urry, W. H. Addition of Carbon Tetrachloride and Chloroform to Olefins. *Science* **1945**, *102*, 128–128.

(18) (a) Matyjaszewski, K.; Xia, J. Atom Transfer Radical Polymerization. *Chem. Rev.* **2001**, *101*, 2921–2990. (b) Eckenhoff, W. T.; Pintauer, T. Copper Catalyzed Atom Transfer Radical Addition (ATRA) and Cyclization (ATRC) Reactions in the Presence of Reducing Agents. *Catal. Rev.: Sci. Eng.* **2010**, *52*, 1–59. (c) Clark, A. J. Copper Catalyzed Atom Transfer Radical Cyclization Reactions. *Eur. J. Org. Chem.* **2016**, *2016*, 2231–2243. (d) Jones, G. R.; Anastasaki, A.; Whitfield, R.; Engelis, N.; Liarou, E.; Haddleton, D. M. Copper-Mediated Reversible Deactivation Radical Polymerization in Aqueous Media. *Angew. Chem., Int. Ed.* **2018**, *57*, 10468–10482. (e) Pan, X.; Fantin, M.; Yuan, F.; Matyjaszewski, K. Externally Controlled Atom Transfer Radical Polymerization. *Chem. Soc. Rev.* **2018**, *47*, 5457–5490.

(19) PPh₃-derived cinchona-alkaloid N,N,P-ligand has been firstly developed by Dixon and coworkers, see: Sladojevich, F.; Trabocchi, A.; Guarna, A.; Dixon, D. J. A New Family of Cinchona-Derived Amino Phosphine Precatalysts: Application to the Highly Enantio- and Diastereoselective Silver-Catalyzed Isocynoacetate Aldol Reaction. *J. Am. Chem. Soc.* **2011**, *133*, 1710–1713.

(20) (a) David, L.; Keith, F. Overview of the Mechanistic Work on the Concerted Metallation–Deprotonation Pathway. *Chem. Lett.* **2010**, *39*, 1118–1126. (b) Davies, D. L.; Donald, S. M. A.; Macgregor, S. A. Computational Study of the Mechanism of Cyclometalation by Palladium Acetate. *J. Am. Chem. Soc.* **2005**, *127*, 13754–13755. (c) García-Cuadrado, D.; Braga, A. A. C.; Maseras, F.; Echavarren, A. M. Proton Abstraction Mechanism for the Palladium-Catalyzed Intramolecular Arylation. *J. Am. Chem. Soc.* **2006**, *128*, 1066–1067. (d) Lafrance, M.; Fagnou, K. Palladium-Catalyzed Benzene Arylation: Incorporation of Catalytic Pivalic Acid as a Proton Shuttle and a Key Element in Catalyst Design. *J. Am. Chem. Soc.* **2006**, *128*, 16496–16497. (e) King, A. E.; Ryland, B. L.; Brunold, T. C.; Stahl, S. S. Kinetic and Spectroscopic Studies of Aerobic Copper(II)-Catalyzed Methoxylation of Arylboronic Esters and Insights into Aryl Transmetalation to Copper(II). *Organometallics* **2012**, *31*, 7948–7957.

(21) Allen, L. A. T.; Raclea, R.-C.; Natho, P.; Parsons, P. J. Recent Advances in the Synthesis of α -Amino Ketones. *Org. Biomol. Chem.* **2021**, *19*, 498–513.

(22) (a) Qi, R.; Wang, C.; Huo, Y.; Chai, H.; Wang, H.; Ma, Z.; Liu, L.; Wang, R.; Xu, Z. Visible Light Induced Cu-Catalyzed Asymmetric C(sp³)-H Alkylation. *J. Am. Chem. Soc.* **2021**, *143*, 12777–12783. (b) Zhou, S.; Zhang, G.; Fu, L.; Chen, P.; Li, Y.; Liu, G. Copper-Catalyzed Asymmetric Cyanation of Alkenes via Carbonyl-Assisted Coupling of Alkyl-Substituted Carbon-Centered Radicals. *Org. Lett.* **2020**, *22*, 6299–6303.